


AMENDMENTS TO THE CLAIMS

1. (Currently Amended) A compound of formula I



wherein

R^1 is selected from hydrogen, C_1 - C_4 alkyl, C_3 - C_6 cycloalkyl, aryl and heteroaryl, wherein the aryl or heteroaryl may be substituted by C_1 - C_4 alkyl;

R^2 is selected from hydrogen and C_1 - C_4 alkyl;

R^3 is selected from hydrogen, C_1 - C_4 alkyl, F, CF_3 , CHF_2 and CH_2F ;

R^4 is selected from hydrogen, F, CF_3 , CHF_2 , CH_2F and CH_3 ;

R^5 is selected from hydrogen and F;

R^5 R^6 is selected from hydrogen and F;

Q is S, NH or NCH_3 , optionally substituted by C_1 - C_4 alkyl;

Y^1 is selected from hydrogen; halogen; nitrile; C_1 - C_4 alkoxy; C_1 - C_4 alkyl wherein one or more of the hydrogen atoms of the alkyl group may be substituted for a fluorine atom;


benzyloxy; nitro in the meta or para position; and C_1 - C_4 alkyl ester;

Y^2 is selected from hydrogen; halogen; nitrile; C_1 - C_4 alkoxy; C_1 - C_4 alkyl wherein one or more of the hydrogen atoms of the alkyl group may be substituted for a fluorine atom; and C_1 - C_4 alkyl ester;

Y^3 is selected from hydrogen; halogen; nitrile; C_1 - C_4 alkoxy; C_1 - C_4 alkyl wherein one or more of the hydrogen atoms of the alkyl group may be substituted for a fluorine atom; and C_1 - C_4 alkyl ester; or

Y^1 and Y^2 may form an aromatic or non-aromatic ring, optionally substituted by halogen, nitrile, C_1 - C_4 alkoxy, C_1 - C_4 alkyl wherein one or more of the hydrogen atoms of the alkyl group may be substituted for a fluorine atom, benzyloxy or C_1 - C_4 alkyl ester; as well as pharmaceutically acceptable salts, hydrates, isoforms and/or optical isomers thereof.

2. (Currently Amended) A compound of formula I



wherein

R^1 is selected from hydrogen, C_1 - C_4 alkyl, C_3 - C_6 cycloalkyl, aryl and heteroaryl, wherein the aryl or heteroaryl may be substituted by C_1 - C_4 alkyl;

R^2 is selected from hydrogen and C_1 - C_4 alkyl;

R^3 is selected from hydrogen, C_1 - C_4 alkyl, F, CF_3 , CHF_2 and CH_2F ;

R^4 is selected from hydrogen, F, CF_3 , CHF_2 , CH_2F and CH_3 ;

R^5 is selected from hydrogen and F;

R^5 R^6 is selected from hydrogen and F;

Q is S, NH or NCH_3 , optionally substituted by C_1 - C_4 alkyl;

Y^1 is selected from hydrogen, halogen, nitrile, C_1 - C_4 alkoxy, and C_1 - C_4 alkyl;

Y^2 is selected from hydrogen, halogen, nitrile, C_1 - C_4 alkoxy, and C_1 - C_4 alkyl;

Y^3 is selected from hydrogen, halogen, nitrile, C_1 - C_4 alkoxy, and C_1 - C_4 alkyl;

as well as pharmaceutically acceptable salts, hydrates, isoforms and/or optical isomers thereof.

3. (Original) A compound according to formula I of claim 1 or 2, wherein

R^1 is hydrogen or C_1 - C_3 alkyl;

R^2 is hydrogen;

R^3 is selected from hydrogen and methyl;

R^4 is hydrogen;

R^5 is hydrogen;

R^6 is hydrogen;

Q is S, NH or NCH_3 , optionally substituted by C_1 - C_4 alkyl;

Y¹ is selected from hydrogen, chloro, C₁-C₂ alkoxy, and C₁-C₂ alkyl; and

Y² is selected from hydrogen, chloro, C₁-C₂ alkoxy, and C₁-C₂ alkyl; and

Y³ is hydrogen.

4. (Original) A compound according to claim 1 selected from *N*-[3-(6-methylpyridin-2-yl)prop-2-yn-1-yl]aniline;
- N*-benzyl-3-(6-methylpyridin-2-yl)prop-2-yn-1-amine;
- N*-methyl-*N*-[3-(6-methylpyridin-2-yl)prop-2-yn-1-yl]aniline;
- (3-methylphenyl)[3-(6-methylpyridin-2-yl)prop-2-yn-1-yl]amine;
- (3-methoxyphenyl)[3-(6-methylpyridin-2-yl)prop-2-yn-1-yl]amine;
- (3-chlorophenyl)[3-(6-methylpyridin-2-yl)prop-2-yn-1-yl]amine;
- [(3-phenylprop-2-yn-1-yl)thio]benzene;
- 1-methoxy-3-[(3-phenylprop-2-yn-1-yl)thio]benzene;
- 2-{3-[(3-chlorophenyl)thio]but-1-yn-1-yl}-6-methylpyridine;
- 2-methyl-6-[3-(phenylthio)prop-1-yn-1-yl]pyridine;
- 2-{3-[(3-chlorophenyl)thio]prop-1-yn-1-yl}-6-methylpyridine;
- 2-{3-[(3-methoxyphenyl)thio]prop-1-yn-1-yl}-6-methylpyridine;
- 2-methyl-6-{3-[(3-methylphenyl)thio]prop-1-yn-1-yl}pyridine;
- (*RS*)-2-{3-[(3-methoxyphenyl)thio]but-1-yn-1-yl}-6-methylpyridine;
- 2-[3-(3-chlorophenyl)-4-methylpent-1-yn-1-yl]-6-methylpyridine;
- 2-{3-[(3,4-dimethylphenyl)thio]prop-1-yn-1-yl}-6-methylpyridine;
- 2-{3-[(3,5-dimethylphenyl)thio]prop-1-yn-1-yl}-6-methylpyridine;

2-{3-[(3-ethoxyphenyl)thio]prop-1-yn-1-yl}-6-methylpyridine;


2-{3-[(4-*tert*-butylphenyl)thio]prop-1-yn-1-yl}-6-methylpyridine; and

2-{3-[(3-chlorophenyl)thio]pent-1-yn-1-yl}-6-methylpyridine.


5. (Currently Amended) A compound according to ~~any one of claims 1-4~~ claim 1 for use in therapy.
6. (Original) A compound according to claim 5, wherein the therapy is treatment or prevention of gastroesophageal reflux disease.

CLAIMS 7-8 (CANCELLED)


9. (Original) A pharmaceutical composition comprising a compound of formula I of claim 1 or 2 as an active ingredient, together with a pharmacologically and pharmaceutically acceptable carrier.
10. (Original) A process for the preparation of a compound of formula I, whereby a coupling reaction of the aryl bromide A




and the alcohol B



is performed in the presence of a base such as triethyl amine, giving the alcohol C



which is then converted into the mesylate D



and reacted with primary or secondary amines or a thiol nucleophile, and wherein

R^1 is selected from hydrogen, C_1 - C_4 alkyl, C_3 - C_6 cycloalkyl, aryl and heteroaryl, wherein

the aryl or heteroaryl may be substituted by C_1 - C_4 alkyl;

R^2 is selected from hydrogen and C_1 - C_4 alkyl;

R^3 is selected from hydrogen, C_1 - C_4 alkyl, F, CF_3 , CHF_2 and CH_2F ;

R^4 is selected from hydrogen, F, CF_3 , CHF_2 , CH_2F and CH_3 ;

R^5 is selected from hydrogen and F;

R^6 is selected from hydrogen and F.

11. (Original) A compound selected from (RS)-4-(6-methylpyridin-2-yl)but-3-yn-2-ol; 4-methyl-1-(6-methylpyridin-2-yl)pent-1-yn-3-ol; Methanesulfonic acid 3-pyridin-2-yl-prop-2-ynyl ester; and 1-(6-Methyl-pyridin-2-yl)-pent-1-yn-3-ol.

12. (Original) A method for the inhibition of transient lower esophageal sphincter relaxations whereby an effective amount of a compound of formula I of claim 1 or 2 is administered to a subject in need of such inhibition.

13. (Original) A method for the treatment or prevention of gastroesophageal reflux disease, whereby an effective amount of a compound of formula I of claim 1 or 2 is administered to a subject in need of such treatment or prevention.